

GUT Precursors and Non-Trivial Fixed Points in Higher-Dimensional Gauge Theories

Keith R. Dienes¹, Emilian Dudas^{2,3}, Tony Gherghetta⁴

¹ *Department of Physics, University of Arizona, Tucson, AZ 85721 USA*

² *LPT, Bât. 210, Univ. Paris-Sud, F-91405, Orsay Cedex, France*

³ *Centre de Physique Théorique, Ecole Polytechnique, F-91128, Palaiseau Cedex, France*

⁴ *School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455 USA*

(October 29, 2002)

Within the context of traditional logarithmic grand unification at $M_{\text{GUT}} \approx 10^{16}$ GeV, we show that it is nevertheless possible to observe certain GUT states such as X and Y gauge bosons at lower scales, perhaps even in the TeV range. We refer to such states as “GUT precursors.” Such states offer an interesting alternative possibility for new physics at the TeV scale, even when the scale of gauge coupling unification remains high, and suggest that it may be possible to probe GUT physics directly even within the context of high-scale gauge coupling unification. More generally, our results also suggest that it is possible to construct self-consistent “hybrid” models containing widely separated energy scales, and give rise to a Kaluza-Klein realization of non-trivial fixed points in higher-dimensional gauge theories. We also discuss how such theories may be deconstructed at high energies.

I. INTRODUCTION

One of the most important theoretical challenges in physics is to determine the nature of fundamental theories. Such fundamental theories include theories of grand unification, quantum gravity, and even strings, with each theory carrying its own intrinsic energy scale.

The traditional view of such theories stipulates that their intrinsic energy scales are exceedingly high. In such cases, experimental evidence in favor of such theories is at best indirect. More recently, however, it has been suggested [1–4] that the presence of large extra dimensions might significantly lower the energy scales associated with such theories, perhaps all the way to the TeV range. In such cases, we might hope for direct experimental tests of such theories.

In this paper, we wish to propose a “hybrid” possibility. Specifically, we wish to consider a higher-dimensional scenario in which the fundamental theories of physics retain their traditional high characteristic energy scales, but in which it is nevertheless possible to obtain *direct*, low-energy evidence of their existence. As we shall see, this will be possible because of the emergence of a non-trivial fixed point which enables a large separation of scales to exist within a single model.

For concreteness, we shall concentrate on the case of grand unification, and consider a scenario in which the unification of gauge couplings retains its traditional logarithmic behavior, with unification occurring near $M_{\text{GUT}} \approx 10^{16}$ GeV. This unification is therefore precisely as in the Minimal Supersymmetric Standard Model (MSSM). However, we shall demonstrate that even within such a scenario, it is possible that certain states associated with the emergence of a grand unified theory

(GUT) at this energy scale can actually be extremely light, perhaps even in the TeV range. We shall refer to such states as “GUT precursors.” The appearance of such precursor states would then provide a direct, experimental window into high-scale, fundamental physics.

II. GUT SYMMETRY BREAKING, ORBIFOLDS, AND “GUT PRECURSORS”

In theories which exhibit a unification of the Standard Model (SM) gauge couplings, it is natural to imagine the emergence of a grand unified theory at the scale of unification. The gauge symmetry group associated with this GUT [*e.g.*, $SU(5)$ or $SO(10)$] must then be large enough to contain the SM gauge symmetry group as a subgroup. In each case, we then find that the corresponding GUT contains not only the usual Standard Model particles, but also additional particles which are directly associated with the GUT. These particles necessarily include the so-called X and Y gauge bosons associated with the enlarged GUT gauge symmetry, and may also include additional matter particles (such as colored Higgs triplets).

There are two basic methods by which GUT symmetries can be broken below the scale of unification. The first method is intrinsically field-theoretic: one imagines that a certain GUT field obtains a non-vanishing expectation value $v \approx M_{\text{GUT}}$ in such a way that the Standard Model fields remain light while the extra GUT fields become heavy. This is the standard Higgs mechanism for breaking a GUT symmetry, and it is characterized by the fact that the masses of the extra GUT fields—the true signatures of the existence of the GUT—are parametrically tied to the GUT scale.

The second method, by contrast, is essentially string-theoretic, and involves truncating the full string Fock space in such a way that the large initial gauge symmetry is broken down to a smaller residual gauge symmetry. This method has a long history in the string literature [5,6], and is the method by which large string gauge symmetries such as $SO(32)$ or $E_8 \times E_8$ are broken down to the Standard Model gauge group in various phenomenological string models [7,8]. This method has often been referred to as “GUT breaking by orbifolds,” and has been discussed within the context of large extra dimensions in Ref. [3] as well as more recently in Refs. [9–12].

This method works as follows. For simplicity, let us first imagine compactification of a single extra dimension on the circle S^1 defined by identifying $y \approx y + 2\pi R$ (where y is the coordinate along the compact extra dimension). Any field $\Phi(x^\mu, y)$ on the circle can then be Fourier-decomposed as $\Phi = \Phi_+ + \Phi_-$ where

$$\begin{aligned}\Phi_+(x^\mu, y) &= \sum_{n=0}^{\infty} \Phi_+^{(n)}(x^\mu) \cos\left(\frac{ny}{R}\right), \\ \Phi_-(x^\mu, y) &= \sum_{n=1}^{\infty} \Phi_-^{(n)}(x^\mu) \sin\left(\frac{ny}{R}\right).\end{aligned}\quad (1)$$

Note that Φ_+ is even under $y \rightarrow -y$, while Φ_- is odd; moreover Φ_- lacks a zero mode. The mass of the Kaluza-Klein mode $\Phi_\pm^{(n)}$ is n/R . Given this Kaluza-Klein circle decomposition, it is then straightforward to compactify on the orbifold defined by S^1/Z_2 where the Z_2 action is $y \rightarrow -y$: we simply retain only the even or odd modes in the above decomposition. It is this truncation which reduces the Fock space. For example, if Φ refers to a Standard Model field, we arrange our orbifold so as to retain the even components Φ_+ , whereas if Φ refers to a GUT field which is not present in the Standard Model, we retain the odd components Φ_- . In this way, the spectrum of zero-modes accessible to the low-energy observer at energies $E \ll R^{-1}$ consists of only the Standard Model fields. In other words, the orbifold projection has broken the GUT at low energies.

For our purposes, however, the important feature of this method of GUT symmetry breaking is the energy scale at which the first signatures of the full GUT symmetry appear. Unlike the Higgs breaking mechanism, where masses of the GUT fields beyond the Standard Model are parametrically tied to M_{GUT} , in this case the masses of the first Kaluza-Klein modes for these GUT particles are set by the inverse radius of the orbifold! Thus, in cases for which $R^{-1} < M_{\text{GUT}}$, we actually begin to observe GUT particles (such as X and Y gauge bosons) *before* we detect actual gauge coupling unification. In other words, these low-lying Kaluza-Klein modes of the GUT particles appear as “GUT precursors” [3], signalling the future emergence of a full gauge coupling unification at an even higher energy scale.

The obvious question, then, is to determine how light these GUT precursors can be. How far below M_{GUT} , the scale of gauge coupling unification, can these states sit?

Clearly, the answer to this question depends on the particular model under discussion. In the models of accelerated power-law unification in Ref. [3], the ratio between the scale of extra dimensions and the scale of accelerated gauge coupling unification is never significantly more than one order of magnitude: $M_{\text{GUT}}R \lesssim 20$. Likewise, in the more recent models of Ref. [10] for which M_{GUT} takes a high value $M_{\text{GUT}} > 10^{16}$ GeV, this ratio is somewhat larger: $M_{\text{GUT}}R \approx 100$. Thus, both classes of models predict the appearance of GUT precursors well in advance of actual gauge coupling unification.

Despite these facts, both classes of models predict the appearance of GUT precursors which are not drastically separated from their corresponding fundamental scales of gauge coupling unification. In other words, neither class of models provides for an extremely large energy range over which the effective theory is higher-dimensional with non-unified gauge couplings.

At first glance, it might appear that we cannot separate these scales too greatly because the GUT precursors themselves will affect the running of the gauge couplings and thereby alter the gauge coupling unification which is responsible for setting the value M_{GUT} . Indeed, the cumulative effects of the Kaluza-Klein excitations of *Standard Model* fields give rise to a power-law running for the gauge couplings which tends to accelerate the scale of gauge coupling unification, thereby tying M_{GUT} closely to the precursor scale R^{-1} .

However, when the effects of the GUT precursors are also included, it then follows that the states at each excited Kaluza-Klein mass level fall into complete GUT multiplets. As originally noticed in Ref. [3], this still leads to gauge coupling unification. However, as pointed out in Ref. [10], the presence of complete GUT multiplets at each excited level implies that the power-law running is *universal* for each gauge coupling. Thus, the *unification* of the gauge couplings continues to be logarithmic, occurring just as it does in four dimensions in the absence of Kaluza-Klein states. Indeed, this feature is the hallmark of the models of Ref. [10], and is one of the reasons why these models can apparently tolerate the larger value $M_{\text{GUT}}R \approx 100$.

The chief danger inherent in such a unification, however, is that the gauge couplings are each individually still experiencing power-law evolution. Thus, even though the *unification* of these gauge couplings is completely logarithmic, the gauge couplings themselves might flow towards strong coupling, thereby invalidating the perturbative unification calculation. Indeed, this is the primary feature which ultimately limits the separation between the scale at which the GUT precursors appear and the scale of gauge coupling unification.

III. POWER-LAW RUNNING AND PERTURBATIVITY

We shall now demonstrate that this restriction does not arise for models in which all matter is restricted to orbifold fixed points and in which only the Standard Model and GUT gauge bosons propagate in the bulk. In such cases, we shall show that it is possible to separate R^{-1} , the scale of the GUT precursors, *by an arbitrary amount* from M_{GUT} , at least as far as gauge coupling unification and perturbativity are concerned. We shall defer our discussion of the general interpretation and phenomenological implications of these results to Sects. IV and VII.

In theories with extra dimensions, the evolution of the gauge couplings takes the approximate form [3]

$$\alpha_i^{-1}(\Lambda) \approx \alpha_i^{-1}(M_Z) - \frac{b_i}{2\pi} \ln \frac{\Lambda}{M_Z} + \frac{\tilde{b}_i}{2\pi} \ln \Lambda R - \frac{\tilde{b}_i X_\delta}{2\pi\delta} [(\Lambda R)^\delta - 1] . \quad (2)$$

The emergence of power-law behavior is expected in a higher-dimensional gauge theory [13,3], and can equivalently be viewed as logarithmic running with a beta-function coefficient that continually changes as successive Kaluza-Klein thresholds are crossed. In Eq. (2), M_Z is our chosen low-energy reference scale; Λ is an arbitrary high scale (ultimately associated with the cutoff of the higher-dimensional gauge theory); δ is the number of compactified dimensions; R is their common radius of compactification; and the normalization factor X_δ is the compactification volume with all radii normalized to unity. For example, for toroidal compactifications we have $X_\delta \equiv \pi^{\delta/2}/\Gamma(1 + \delta/2)$ (the volume of a δ -dimensional unit sphere). Likewise, b_i are the beta-function coefficients of the zero-mode fields (including the contributions of those fields which do not feel extra dimensions altogether, such as those restricted to branes and/or orbifold fixed points), while $\tilde{b}_i$ are the beta-function coefficients associated with the field content at each excited Kaluza-Klein level. Thus, it is the latter beta-function coefficients which govern the intrinsically higher-dimensional power-law evolution of the gauge couplings. Note that in the case of compactification on an orbifold, the $\tilde{b}_i$ -dependent contributions in Eq. (2) are generally reduced due to the orbifold projections at the excited Kaluza-Klein levels; with a $\mathbb{Z}_2$ projection, for example, these terms must be divided by 2. Also note that if the zero-mode fields are those of the MSSM, then the sum of the first two terms on the right side of Eq. (2) is equal to the usual MSSM value of the (inverse) unified gauge coupling when $\Lambda = M_{\text{GUT}} \approx 2 \times 10^{16}$ GeV.

In cases where the $\tilde{b}_i$ are unequal, we see from Eq. (2) that the power-law evolution of the gauge couplings is different for each gauge coupling. This implies that the relative *differences* between the gauge couplings also evolve

with power-law behavior. However, when the $\tilde{b}_i$ are all equal, we see from Eq. (2) that this power-law behavior is universal for all gauge couplings. The relative differences of gauge couplings then evolve purely logarithmically, exactly as in four dimensions. Indeed, even the $\tilde{b}_i$ -dependent logarithmic contributions in Eq. (2) are universal, and do not affect the unification of the couplings. Thus, we find that when the $\tilde{b}_i$ are all equal, we retain exactly the same logarithmic gauge coupling unification that arises in four dimensions.

Despite this fact, it is still important to verify that the individual gauge couplings themselves remain perturbative over the entire energy range from R^{-1} to $\Lambda \equiv M_{\text{GUT}}$. Otherwise, the use of the one-loop result in Eq. (2) is no longer valid. Towards this end, let us assume that $\tilde{b}_i \equiv \tilde{b} < 0$ for all i . Since $\tilde{b} < 0$, the power-law contributions to the gauge couplings push the couplings towards extremely weak values. Indeed, in the limit where $\Lambda R \gg 1$, we find from Eq. (2) that each of the gauge couplings scales in the ultraviolet as

$$\alpha(\Lambda) \approx -\frac{2\pi\delta}{\tilde{b}X_\delta}(\Lambda R)^{-\delta} . \quad (3)$$

However, even though these couplings are extremely weak, the true loop expansion parameter in such a situation is $\alpha_{\text{eff}} \equiv N\alpha$ where $N \equiv X_\delta(\Lambda R)^\delta$ is the number of Kaluza-Klein levels that have been crossed. Indeed, α_{eff} describes the effective strength of the gauge interaction, since it characterizes the coupling of each individual Kaluza-Klein mode multiplied by the multiplicity of these modes. Thus, for true perturbativity, we must demand $\alpha_{\text{eff}} \ll 4\pi$.

Remarkably, this constraint is satisfied no matter how large ΛR becomes. Indeed, we find that $\alpha_{\text{eff}} \approx -2\pi\delta/\tilde{b}$ as $\Lambda R \rightarrow \infty$, so that the condition for perturbativity becomes $-\delta/(2\tilde{b}) \ll 1$. Thus, as long as $\tilde{b}$ is sufficiently large and negative, this condition can be satisfied even if $\Lambda R \gg 1$.

As an example, let us consider a scenario in which, as discussed above, the zero-mode fields are those of the MSSM and only the GUT gauge bosons sit in the bulk. For simplicity, we shall take our unified gauge group to be $SU(5)$, and we shall also assume that $\delta = 1$. Since our low-energy theory is $\mathcal{N} = 1$ supersymmetric, the bulk fields necessarily fall into $\mathcal{N} = 2$ supermultiplets. Our bulk fields therefore consist of $\mathcal{N} = 2$ vector multiplets transforming in the adjoint of $SU(5)$, leading to $\tilde{b}_i = \tilde{b} = -10$ for all i . We then find that the effective gauge interaction strength at unification is $\alpha_{\text{eff}} \approx 0.63$, which is considerably less than 4π . Note that this remains true even if $\Lambda R \approx 10^{13}$. Thus it is possible for the GUT precursors to appear at the TeV scale even though the (logarithmic) gauge coupling unification does not occur until the usual scale $M_{\text{GUT}} \approx 10^{16}$ GeV.

This behavior is illustrated in Fig. 1, where we plot the value of the effective unified coupling α_{eff} at M_{GUT}




FIG. 1. The effective unified coupling $\alpha_{\text{eff}}(M_{\text{GUT}})$ as a function of $M_{\text{GUT}}R$ for the five-dimensional $SU(5)$ GUT model discussed in the text. This coupling remains perturbative for arbitrarily large values of $M_{\text{GUT}}R$.

as a function of R , holding M_{GUT} fixed at its usual four-dimensional value 2×10^{16} GeV. We have taken $\delta = 1$ and $\tilde{b} = -10$, as discussed above. It is clear that the effective coupling remains perturbative for arbitrarily large values of $M_{\text{GUT}}R$, saturating at its asymptotic value as early as $M_{\text{GUT}}R \approx 100$. Thus, the scale at which our GUT precursors appear can be separated by an arbitrary amount from the scale at which the gauge couplings unify.

Note that this scenario requires beta-function coefficients $\tilde{b}_i$ which are universal and *negative*. Ordinarily, it might seem impossible to arrange $\tilde{b}_i$ negative for each gauge group simultaneously, since in most scenarios (such as the MSSM) the hypercharge coupling typically becomes stronger with increasing energy. However, $\tilde{b}_i$ is negative in our scenario precisely as a result of the GUT precursors. Indeed, in order to obtain a negative hypercharge beta-function coefficient, the X and Y gauge bosons are just what we require: spacetime vector bosons with non-trivial hypercharge assignments. Thus, it is the GUT precursors which permit the theory to remain perturbative even as $\Lambda R \gg 1$.

One might worry that two-loop effects might be significant in such a scenario. However, two-loop effects essentially *vanish* in the $\Lambda R \rightarrow \infty$ limit, since the presence of $\mathcal{N} = 2$ supersymmetry in the bulk ensures that the higher-loop power-law effects are suppressed by a factor of $1/\Lambda R$ relative to the one-loop effects [3,15]. Even when ΛR remains finite, it is straightforward to verify that two- and higher-loop corrections do not substantially alter the logarithmic unification which emerges at

one-loop order [3,15]. Likewise, we remark in this context that the potentially damaging brane surface kinetic terms discussed in Ref. [16] become vanishingly small in this context, since the volume of the bulk becomes 10^{13} as large as the volume on the brane. Thus, the unification of the gauge couplings is maintained.

It is also important to understand what happens if $\tilde{b} > 0$. In this case, the gauge couplings become stronger rather than weaker as we evolve upwards in energy, ultimately hitting a Landau pole where $\alpha^{-1}(\Lambda) = 0$. We emphasize that this Landau pole is completely physical, since it is induced by the cumulative effects of the Kaluza-Klein thresholds as we evolve upwards in energy. [Even if we interpret Eq. (2) as a threshold correction rather than as a renormalization group “running,” when $\tilde{b} > 0$ this “correction” can push α^{-1} to negative values, which is clearly unacceptable.] In such cases, depending on the specific value of $\tilde{b}$, there is therefore a maximum allowed value of $M_{\text{GUT}}R$ which can be tolerated if we imagine varying R while holding M_{GUT} fixed. For example, in some of the “minimal” models of Ref. [10], the bulk contains not only the $SU(5)$ gauge bosons, but also two **5** representations (for the Higgses) and four **10** representations (for the first two generations). Since these are all $\mathcal{N} = 2$ supermultiplets, this leads to a value $\tilde{b} = +4$. (In general, with n_5 multiplets transforming in the **5** representation and n_{10} multiplets in the **10** representation, one obtains $\tilde{b} = -10 + n_5 + 3n_{10}$.) We then find that perturbativity in such models requires a maximum value $M_{\text{GUT}}R \lesssim 44$ in order to avoid a Landau pole (or equivalently, to avoid negative inverse gauge couplings). Since some of the models in Ref. [10] require $M_{\text{GUT}}R \approx 100$, it is unclear how the perturbative logarithmic unification prediction is maintained in such models.

IV. POWER-LAW RUNNING AND NON-TRIVIAL ULTRAVIOLET FIXED POINTS

In the setup described in the previous section, the asymptotic ultraviolet power-law scaling of gauge couplings towards weak values is exactly compensated by the asymptotic power-law growth of the number of degrees of freedom in the theory in such a way that the product of these two quantities remains a constant. Thus, the effective strength of the gauge interactions appears to approach a non-trivial fixed point in the ultraviolet. Such behavior for gauge couplings with $\tilde{b} < 0$ was also observed previously in Ref. [14]. Of course, in our particular scenario we do not need to approach the truly asymptotic limit $\Lambda R \rightarrow \infty$ because the logarithmic terms in Eq. (2) eventually induce a unification at $\Lambda \equiv M_{\text{GUT}}$; we therefore can take this as a cutoff for the higher-dimensional running. Nevertheless, it is intriguing to consider the formal limit in which we disregard issues of unification and take $\Lambda R \rightarrow \infty$.

It is already apparent from Fig. 1 that as long as $\Lambda R \gtrsim 100$, our theory essentially becomes “scale invariant” in the sense that the ultraviolet physics becomes independent of the low-energy scale R^{-1} at which the GUT precursors appear. We may also rephrase this observation directly in terms of the effective couplings $\alpha_{\text{eff},i} \equiv N\alpha_i$ where $N \equiv X_\delta(\Lambda R)^\delta$. Given the evolution equations for the couplings α_i in Eq. (2), it is straightforward to show that the effective couplings $\alpha_{\text{eff},i}$ evolve according to

$$\Lambda \frac{d\alpha_{\text{eff},i}^{-1}}{d\Lambda} = - \left(\delta\alpha_{\text{eff},i}^{-1} + \frac{\tilde{b}_i}{2\pi} \right) + \left(\frac{\tilde{b}_i - b_i}{2\pi X_\delta} \right) (\Lambda R)^{-\delta} + \frac{c_i}{2\pi} \frac{\alpha_{\text{eff},i}}{4\pi} (\Lambda R)^{-\delta} + \dots, \quad (4)$$

where in the second line we have written the dominant two-loop contributions arising from the bulk and boundary fields running in the loops (with c_i representing a two-loop beta-function coefficient). Thus, even though the individual gauge couplings α_i themselves evolve with power-law behavior, we see from Eq. (4) that for $\Lambda R \gg 1$, the *effective* gauge couplings $\alpha_{\text{eff},i}$ each approach an ultraviolet fixed point at $\alpha_{\text{eff},i} = -2\pi\delta/\tilde{b}_i$. Moreover, if $\tilde{b}_i \equiv \tilde{b}$ for all i , we see that even though the differences of the gauge couplings continue to evolve logarithmically, the fixed-point values of the *effective* gauge couplings all become equal. Thus, in this sense, we see that the effective strengths of the gauge interactions in this theory each flow to a *common* fixed point in the ultraviolet. Note that two- and higher-loop effects merely contribute additional power-law terms in Eq. (4) which again vanish in the $\Lambda R \rightarrow \infty$ limit. Such contributions therefore do not alter the ultraviolet fixed-point structure of these theories.

It is natural to interpret these results as indicating the emergence of a non-trivial (interacting) ultraviolet fixed point corresponding to a supersymmetric, higher-dimensional, unified gauge theory. Indeed, such higher-dimensional fixed-point gauge theories are known to exist in uncompactified five [17] and six [18] dimensions. Since we expect the ultraviolet (short-distance) limit of our compactified theory to reproduce the physics of an uncompactified higher-dimensional theory, it is tempting to identify the ultraviolet limit of our theory as one of the interacting fixed-point theories discussed in Refs. [17,18].

As an example, let us consider the case of $SU(N)$ gauge theory in five dimensions. If the only matter consists of n_f “quarks” transforming in the fundamental representation, then the necessary and sufficient condition [17] for the existence of an interacting ultraviolet fixed point is $n_f \leq 2N$. This is equivalent to our requirement that $\tilde{b} \leq 0$.

It is important to stress that for unitary groups, the conditions for the existence of such non-trivial fixed points are generally *stronger* than merely demanding

$\tilde{b} < 0$. For example, if we also introduce bulk matter which transforms in the *antisymmetric tensor* representation [*e.g.*, the **10** representation of $SU(5)$], then further restrictions on the number of fundamental representations must be imposed in order to guarantee the fixed-point behavior of the theory in the ultraviolet [17]. These conditions are consistent with $\tilde{b} \leq 0$, but provide further constraints on the precise matter content. However, our main point is that we can always ensure that our theory flows to a non-trivial fixed point in the ultraviolet by choosing the bulk field content appropriately. This then guarantees the self-consistency of our scenario and the large separation in energy scales that it contains.

One important by-product of this analysis is that it essentially furnishes us with an alternative, four-dimensional “Kaluza-Klein” realization of these fixed-point theories. In such a realization, the effective higher-dimensional gauge coupling at the fixed point asymptotically emerges in the ultraviolet as the product $\alpha_{\text{eff}} = N\alpha$. [Note that the *dimensionful* gauge coupling in higher dimensions is $\alpha_{4+\delta}(\Lambda) = \Lambda^{-\delta}\alpha_{\text{eff}}(\Lambda)$.] Moreover, the existence of such “Kaluza-Klein” realizations for these higher-dimensional fixed points may provide a new tool for studying the properties of these fixed points under compactification, orbifold projection, GUT symmetry breaking, and even supersymmetry breaking. These are precisely the issues that must be addressed if such higher-dimensional fixed-point gauge theories are to have phenomenologically relevant compactifications to four dimensions. Thus, these four-dimensional “Kaluza-Klein” realizations of these interacting fixed points may provide a new approach which can be used when studying how these theories behave under phenomenologically relevant compactifications, and when calculating the subleading corrections that such compactifications introduce.

V. DECONSTRUCTION AND UV/IR FIXED-POINT MATCHING

Recently, a new four-dimensional ultraviolet completion of Kaluza-Klein theories was proposed [19]. When generalized to a supersymmetric context [20], this “deconstruction” proposal embeds the Kaluza-Klein theory into a four-dimensional asymptotically free theory based on the gauge group $SU(N_c)^{N+1}$. The model can also be described by a “moose” (or quiver) diagram with $N+1$ sites and links. At each site on the moose diagram, there is an $\mathcal{N} = 1$ supersymmetric $SU(N_c)$ gauge theory; likewise, the links represent chiral multiplets Φ_I which transform in the bifundamental representation of the two adjacent $SU(N_c)$ gauge groups. Each site therefore contains $N_f = N_c$ flavors transforming in the fundamental representation of the $SU(N_c)_i$ gauge group. At the deconstruction scale, the scalar components ϕ_I within the chiral supermultiplets Φ_I each accrue equal vacuum ex-

pectation values $\langle \phi_I \rangle = v$ which in turn break the gauge group to the diagonal: $SU(N_c)^{N+1} \rightarrow SU(N_c)$. In the energy range $gv/N < \mu < gv$, the resulting spectrum is then similar to that of a pure $\mathcal{N} = 2$ supersymmetric $SU(N_c)$ Yang-Mills theory compactified on a circle.

Deconstruction naturally provides a renormalizable context for analyzing the ultraviolet fixed points discussed in this paper. Note, first of all, that the effective coupling α_{eff} which approaches the ultraviolet fixed-point value represents in the deconstruction scheme the gauge coupling of gauge groups at each site at the deconstruction scale. Moreover, the fact that α_{eff} is independent of N is natural from the deconstruction viewpoint.

However, when embedding these ultraviolet fixed points into a deconstructed theory at even higher energy scales, it is important to maintain the ultraviolet insensitivity of low-energy observables. Otherwise, the underlying fixed-point structure will be destroyed by the deconstruction. Thus, in order to preserve this ultraviolet insensitivity, we see that the ultraviolet fixed points from the Kaluza-Klein perspective must somehow be identified with (or matched with) *infrared* fixed points of the theory above the deconstruction scale.

The obvious candidates for these infrared fixed points are the four-dimensional fixed points proposed in connection with the conjectured $\mathcal{N} = 1$ Seiberg duality [21]. For $SU(N_c)$ gauge groups, it was argued in Ref. [21] that infrared fixed points exist only when the number N_f of flavors satisfies $3N_c/2 < N_f < 3N_c$. Thus, in building our deconstructed theory, we are led to introduce an additional $n_f > N_c/2$ flavors of chiral multiplets $Q_i, \tilde{Q}_i$ at each site. The total number of flavors at each site is therefore $N_f = N_c + n_f$. For our purposes we will not need a full superconformal field theory in the infrared limit of our deconstructed theory, and it suffices merely to impose the existence of an infrared fixed point for the gauge couplings. In our case, the exact beta-function becomes [22]

$$\beta(g) = \frac{g^3}{16\pi^2} \frac{b'}{1 - N_c g^2 / 8\pi^2} \quad (5)$$

where the coefficient b' is given by

$$b' = -2N_c + n_f - \frac{N_c}{2}(\gamma_\Phi^{I-1} + \gamma_\Phi^I) - \frac{n_f}{2}(\gamma_Q^I + \gamma_{\tilde{Q}}^I) \quad (6)$$

and where the one-loop anomalous dimensions of the fields at each site/link I are given by

$$\begin{aligned} \gamma_\Phi &= -\frac{1}{8\pi^2} \left(2 \frac{N_c^2 - 1}{N_c} g^2 - n_f \lambda^2 \right), \\ \gamma_Q &= \gamma_{\tilde{Q}} = -\frac{1}{8\pi^2} \left(\frac{N_c^2 - 1}{N_c} g^2 - N_c \lambda^2 \right). \end{aligned} \quad (7)$$

The superpotential describing the deconstructed theory is given by

$$W = \sum_{i,I} (\sqrt{2} \lambda \tilde{Q}_{i,I} \Phi_I Q_{i,I+1} - m \tilde{Q}_{i,I} Q_{i,I}), \quad (8)$$

whereupon we see that the Kaluza-Klein spectrum for the n_f hypermultiplets is recovered by setting $m = \sqrt{2} g^2 v / \lambda$. In addition, the zero mode of the hypermultiplets has a mass $m_0^2 = 2v^2(g^2 - \lambda^2)^2 / \lambda^2$. Since no site is singled out, we have taken $\gamma_\Phi^I \equiv \gamma_\Phi$ for all I in Eq. (7). We then find that requiring $\beta(g) = 0$ defines a curve of fixed points in the (g, λ) plane.

Let us now consider how to match the ultraviolet fixed point of the five-dimensional theory with the infrared fixed point of the deconstructed theory. First, we must take $\lambda = g$ in our deconstructed theory in order to guarantee that the interactions below the deconstruction scale respect $\mathcal{N} = 2$ supersymmetry. Second, since our theory below the deconstruction scale has $SU(N_c)$ gauge symmetry with n_f flavors, we see that $\tilde{b} = -2N_c + n_f$. Thus, our Kaluza-Klein theory has $g_{\text{eff}}^2 = 8\pi^2 / (2N_c - n_f)$.

In order to match the ultraviolet and infrared fixed points, we now require solutions for $\beta(g) = 0$ to occur at $g = g_{\text{eff}}$. Setting $b' = 0$ and requiring $g = g_{\text{eff}}$, we obtain the constraint

$$(2N_c - n_f)(n_f - N_c) = 2 + \frac{n_f}{N_c}. \quad (9)$$

In the large- N_c limit, this has two solutions: $n_f = N_c$ and $n_f = 2N_c$. We can reject the second solution because it implies $\tilde{b} = 0$, or $g_{\text{eff}} \rightarrow \infty$. (Equivalently, taking $n_f = 2N_c$ violates the infrared fixed-point constraint $N_f < 3N_c$ in the original deconstructed theory.) However, the solution with $n_f = N_c$ yields $N_c \alpha_{\text{eff}} = 2\pi$, which is finite and semi-perturbative. Of course, we must be slightly more careful because this value for g_{eff} causes the denominator in Eq. (5) to diverge. However, for any large but finite value of N_c , there exists an exact solution for n_f/N_c in Eq. (9) which is only slightly larger than 1. For this exact solution, we find that $b' = 0$ and consequently $\beta(g) = 0$. Note that higher-loop contributions in Eq. (7) are suppressed by the small gauge coupling and thus provide only a small correction to the exact solution for n_f/N_c in Eq. (9).

Of course, a fully non-perturbative solution for the ultraviolet/infrared fixed-point matching might also exist. In particular, if we demand that the infrared limit of our deconstructed theory is actually superconformal, then it may be possible to achieve non-perturbative solutions to the constraint $b' = 0$ if the full, non-perturbative, anomalous dimensions satisfy

$$\gamma_\Phi = -2, \quad \gamma_Q + \gamma_{\tilde{Q}} = 2. \quad (10)$$

The non-perturbative solution to these equations should therefore match the ultraviolet fixed point of our Kaluza-Klein theory in order to preserve the ultraviolet insensitivity of the low-energy theory.

Thus, we conclude that it is possible, in principle, to deconstruct our ultraviolet fixed points while maintaining their ultraviolet insensitivity. Note that the embedding into a deconstructed theory also ensures that there are no other non-renormalizable operators which could appear in the ultraviolet theory and cause difficulties when extrapolated down to lower energies.

VI. ORBIFOLD CHOICES AND NON-UNIVERSAL LOGARITHMS

Thus far, we have presented a general GUT scenario in which GUT precursor states can be extremely light compared with the scale of gauge coupling unification. Our purpose has been to illustrate the emergence and utilization of non-trivial fixed points as a means of incorporating widely separated energy scales within a single model. However, in order to build a fully consistent model, we must choose an explicit orbifold and take into account certain additional contributions that arise [10,23–25].

Let us first consider the case of the five-dimensional $SU(5)$ theory compactified on the $S_1/\mathbb{Z}_2$ orbifold presented in Sect. II. Under the $\mathbb{Z}_2$ action, Standard-Model gauge supermultiplets are even while GUT supermultiplets such as the X and Y gauge supermultiplets are odd. However, since this is a five-dimensional theory, the X and Y supermultiplets also have fifth components X_5 and Y_5 , and consistency of the $\mathbb{Z}_2$ orbifold requires that these components be even. The zero modes of such fields then produce additional non-universal logarithmic contributions to the runnings of the gauge couplings, with coefficients $(b_1, b_2, b_3) = (5, 3, 2)$. However, it is easy to verify that taking $R^{-1} \sim \text{TeV}$ still leads to an approximate unification at $M_{\text{GUT}} \approx 2 \times 10^{13} \text{ GeV}$ which, although not as precise as the MSSM unification, is nevertheless more precise than the unification in the Standard Model. Remarkably, as we shall discuss in Sect. VII, such an extra dimension also lowers the fundamental Planck scale to this new unification scale. Of course, at a phenomenological level, it still remains necessary to find a mechanism to give masses to these X_5 and Y_5 zero modes, since they represent colored and fractionally charged scalars.

Another option is to compactify our theory on an $S_1/(\mathbb{Z}_2 \times \mathbb{Z}'_2)$ orbifold with two distinct $\mathbb{Z}_2$ discrete actions [9,10] associated with $y \rightarrow -y$ and $y \rightarrow \pi R - y$. Under the $(\mathbb{Z}_2, \mathbb{Z}'_2)$ orbifold actions, the Standard-Model gauge fields A_μ have $(+, +)$ eigenvalues [with corresponding cosine modings as in Eq. (1) with $n \in 2\mathbb{Z}$ only], while A_5 has $(-, -)$ eigenvalues (resulting in sine modings with $n \in 2\mathbb{Z}$). Likewise, the GUT precursors X_μ and Y_μ have $(-, +)$ eigenvalues (resulting in sine modings with $n \in 2\mathbb{Z} + 1$), while X_5 and Y_5 have $(+, -)$ eigenvalues (resulting in cosine modings with $n \in 2\mathbb{Z} + 1$). With this orbifold choice, only the Standard Model fields have zero modes, but this occurs at the expense of split-

ting the complete GUT multiplets at each Kaluza-Klein level into a subset at even levels, with beta-function coefficients $(\tilde{b}_1, \tilde{b}_2, \tilde{b}_3) = (0, -4, -6)$, and a subset at odd levels, with $(\tilde{b}'_1, \tilde{b}'_2, \tilde{b}'_3) = (-10, -6, -4)$. This results in a staggered, “zig-zag” running for the gauge couplings which averages to a universal power-law running with an effective radius $R/2$, along with a non-universal logarithmic correction. Specifically, the gauge couplings now run according to

$$\alpha_i^{-1}(\Lambda) \approx \alpha_i^{-1}(M_Z) - \frac{b_i}{2\pi} \ln \frac{\Lambda}{M_Z} + \frac{\tilde{b}}{4\pi} \ln \frac{\Lambda R}{2} - \frac{\tilde{b}}{2\pi} \left[\left(\frac{\Lambda R}{2} \right) - 1 \right] - \frac{\tilde{b}'_i}{2\pi} Y \quad (11)$$

where $\tilde{b} \equiv \tilde{b}_i + \tilde{b}'_i = -10$ for all i , where we have neglected various universal additive constants, and where the non-universal logarithm is given by

$$Y \equiv \sum_{n=0}^{(\Lambda R - 2)/2} \ln \frac{2n + 2}{2n + 1} \approx \frac{1}{2} \ln \frac{\pi \Lambda R}{2} \quad (12)$$

with the last approximation holding in the $\Lambda R \gg 1$ limit. Given this running, we then find that the three gauge couplings continue to experience an approximate unification. With $R^{-1} \sim \text{TeV}$, the unification scale is unfortunately quite high ($M_{\text{GUT}} \approx 10^{21} \text{ GeV}$), but increasing R^{-1} not only improves the accuracy of the resulting unification but also lowers the unification scale. Asymptotically, with $R^{-1} \approx 10^{15} \text{ GeV}$, we obtain an essentially *exact* unification at $M_{\text{GUT}} \approx 10^{17} \text{ GeV}$. Although this resembles the energy scales in the scenario in Ref. [10], we stress that the bulk theory here contains only gauge fields, and our gauge couplings become *weak* rather than strong in the ultraviolet limit. Of course, one can also place additional matter in the bulk or on the brane so as to increase the accuracy of the unification and lower the unification scale [25].

A final option is to place a single Higgs five-plet in the bulk. After compactifying on the $S_1/(\mathbb{Z}_2 \times \mathbb{Z}'_2)$ orbifold, the bulk Higgs doublets (triplets) are at even (odd) Kaluza-Klein levels. We thus have $\tilde{b} = -9$, $(\tilde{b}_1, \tilde{b}_2, \tilde{b}_3) = (3/5, -3, -6)$, and $(\tilde{b}'_1, \tilde{b}'_2, \tilde{b}'_3) = (-48/5, -6, -3)$. Note that these values of $\tilde{b}_i$ are the same as those of Ref. [3]. This yields a unification of gauge couplings which is similar to (and approximately as accurate as) the case with only gauge bulk fields discussed above; in each case, small (few-percent) threshold corrections at the unification scale are sufficient to render the unification exact for all values of R^{-1} . Of course, in each case one must place the remaining Higgs field(s) on a Standard-Model brane lacking the GUT symmetry in order to avoid the doublet/triplet problem.

VII. DISCUSSION

In this paper, we have shown that it is possible for certain “GUT precursor” states to appear with masses that are significantly below the scale at which grand unification occurs. As we have seen, the fixed-point structure which makes this possible can be viewed as a general tool which permits the construction of generalized “hybrid” models in which widely separated energy scales can co-exist in a natural way.

Needless to say, these observations prompt a number of important questions, both phenomenological and theoretical. Among the most important phenomenological questions is the issue of proton decay. Ordinarily, light X and Y gauge boson precursors will mediate rapid proton decay. However, as in all low-scale extensions to the Standard Model, this problem may be cured through the use of split fermions on the branes [26] or through the introduction of extra discrete symmetries [3,27]. Likewise, other phenomenological issues include doublet/triplet splitting and general issues of flavor physics. Although we have not attempted to make a complete GUT model that accommodates these phenomena, one could imagine doing so following the lines of Refs. [9–12] except that we now have the interesting option of extending the energy scales of such models into the TeV range. Moreover, we are also free to introduce further matter into the bulk beyond what we have discussed here, provided we ensure that $\tilde{b}$ remains sufficiently large and negative and provided the conditions [17] for an ultraviolet fixed point are maintained. Indeed, the presence of two widely separated scales in such models suggests that we might even try to use the GUT precursors to trigger electroweak symmetry breaking and/or supersymmetry breaking. Of course, we have not speculated on the unknown dynamics which might ultimately be responsible for generating and stabilizing such a large radius; ideas along these lines can be found, *e.g.*, in Ref. [28].

Our results in this paper also raise a number of theoretical issues. The most important concerns the ultraviolet limit of our theory. Although we have shown that the evolution of the gauge couplings is consistent with perturbativity even when the effective higher-dimensional energy interval is large, one must actually verify that *all* correlation functions in the theory remain finite and under control over this large energy range. This is clearly connected with the over-riding question discussed in Sect. IV concerning the manner in which we approach a scale-invariant fixed point in the ultraviolet. By counting Kaluza-Klein states and vertex factors in diagrams with arbitrary numbers of loops and external legs, it is straightforward to demonstrate that all diagrams in this theory necessarily scale as $(N\alpha)^k \sqrt{\alpha}^\ell$ where k and ℓ are non-negative integers. Thus, in the ultraviolet limit, such

diagrams either vanish (if $\ell \neq 0$) or approach a fixed finite value (if $\ell = 0$). Indeed, even though the number of states in this theory is diverging at higher energies, the individual gauge couplings are falling to zero in an exactly compensatory manner.

For example, the four-fermion amplitude for tree-level Kaluza-Klein exchange in the $\delta = 1$ case becomes

$$\begin{aligned} A(s) &\sim \alpha(s) \sum_n \frac{1}{s - n^2/R^2} \sim \frac{\alpha(s)}{\sqrt{s}} R \cot(\pi R \sqrt{s}) \\ &\longrightarrow \frac{\alpha_{\text{eff}}}{s} \cot(\pi R \sqrt{s}) \end{aligned} \quad (13)$$

where we have taken the limit $sR^2 \gg 1$ and identified $\alpha_{\text{eff}} \sim (R\sqrt{s})\alpha(s)$ as $s \rightarrow \infty$. Thus, since $A(s)$ continues to have the asymptotic energy dependence $\sim 1/s$, no unitarity bounds are violated in the ultraviolet. (The divergence when $R\sqrt{s} \in \mathbb{Z}$ merely reflects a fine-tuned Kaluza-Klein resonant pole.) Moreover, as we have already noted, the presence of $\mathcal{N} = 2$ supersymmetry in the bulk ensures the vanishing of many diagrams which would otherwise lead to large and potentially uncontrollable corrections. It will therefore be interesting to explore the ultraviolet self-consistency of this theory further [29]. We stress, however, that amplitudes such as those in Eq. (13) will remain consistent with unitarity bounds only as long as the growing number of Kaluza-Klein states is offset by the power-law running of the gauge couplings. If there are more than two extra dimensions, the power-law running for the gauge couplings will be cancelled by the $\mathcal{N} = 4$ supersymmetry that would be required in the bulk. In such cases, the theory will remain unitary only up to the scale at which these extra dimensions become apparent.

Note that the presence of a large, scale-invariant energy interval in our model is reminiscent of previous models [30] in which “conformality” is invoked to solve the technical hierarchy problem. These models also share certain features with GUT orbifold models formulated in warped backgrounds [31], as well as with string models in which physics at all scales conspires to eliminate the quantum-mechanical sensitivity between light and heavy energy scales [32].

Another important theoretical issue for our models concerns gravity. In this paper we have merely proposed a model of gauge interactions, and as such, any value of R for the GUT precursors is permitted. However, if we wish to incorporate gravity, further complications arise. Since our large extra dimension is presumably also felt by gravity, Kaluza-Klein gravitons will induce Newton’s constant to run more quickly. The standard Gauss-law arguments of Ref. [2] then imply that taking $R^{-1} \sim \mathcal{O}(\text{TeV})$ lowers the effective higher-dimensional Planck scale M_* to approximately 10^{14} GeV. Although this is comfortably within all experimental constraints, this value is slightly below $M_{\text{GUT}} \approx 10^{16}$ GeV. This indicates that we reach a region of strong gravity *before* our gauge couplings unify.

There are two ways in which this situation might be avoided. Within the context of the $S_1/\mathbb{Z}_2$ orbifold discussed in Sect. VI, we have seen that the additional non-universal logarithmic contributions actually lower the unification scale to approximately 10^{13} GeV. Thus, for the $S_1/\mathbb{Z}_2$ orbifold, there is no difficulty. On the other hand, for the $S_1/(\mathbb{Z}_2 \times \mathbb{Z}_2')$ orbifold, we have seen in Sect. VI that the unification scale generally exceeds 10^{16} GeV. One possible solution is then to restrict the GUT precursor scale to the range $R^{-1} \gtrsim 10^{10}$ GeV; this ensures that $M_* \gtrsim M_{\text{GUT}}$. Such GUT precursors would still be significantly lighter than the unification scale, although no longer accessible to upcoming collider experiments. Another solution, however, is to lower the value of M_{GUT} to 10^{14} GeV by introducing further states with appropriate gauge quantum numbers into the theory. Such states would not spoil the universal power-law behavior of this theory if they are restricted to the orbifold fixed points. Gauge coupling unification in such a scenario would then continue to be logarithmic, as desired.

A similar issue arises if we attempt to embed this scenario into string theory. Let us first consider the case of the perturbative heterotic string. In general, the heterotic string scale is related to the Planck scale through the unified gauge coupling: $M_{\text{string}} = g_{\text{GUT}} M_{\text{Planck}}$. However, as a result of the asymptotic power-law running of the gauge couplings, we know that $\alpha_{\text{GUT}}^{-1} \sim (M_{\text{string}} R)^\delta$. (In writing this relation, we are imagining an ultimate identification of M_{string} with M_{GUT} .) However, combining these results and neglecting all numerical factors, we find that $M_{\text{string}} \sim M_*$ where $M_*^{2+\delta} \sim M_{\text{Planck}}^2/R^\delta$ is the higher-dimensional Planck scale at which gravitational effects become significant. Thus, for the heterotic string, the power-law scaling of the gauge coupling implies that it is not possible to separate the scales at which both gravitational and string-theoretic effects become significant. Note that this result persists regardless of the value of the underlying (perturbative) string coupling.

Within the context of Type I strings, however, this situation is somewhat different. For Type I strings, the relation between the string and Planck scales is modified: $M_{\text{string}} = g_{\text{GUT}} \sqrt{g_{\text{string}}} M_{\text{Planck}}$, where g_{string} is the underlying string coupling. Again taking $\alpha_{\text{GUT}}^{-1} \sim (M_{\text{string}} R)^\delta$, we now find the relation

$$M_{\text{string}}^{2+\delta} \sim g_{\text{string}} M_*^{2+\delta}. \quad (14)$$

Thus, since $M_* \lesssim 10^{14}$ GeV (with the maximum value for M_* occurring for $\delta = 1$), we see that we can indeed take $M_{\text{string}} \approx M_{\text{GUT}}$ if we have a large string coupling $g_{\text{string}} \gg 1$. This suggests that embedding our scenario within the context of Type I strings necessarily involves strong couplings and non-perturbative physics. Moreover, since the Type I string coupling is given by $g_{\text{string}} \sim \alpha_{\text{GUT}} V_6 M_{\text{string}}^6$ where V_6 is the *to-*

tal six-dimensional compactification volume, we see that taking $g_{\text{string}} \gg 1$ requires $M_{\text{string}}^6 V_6 \gg M_{\text{string}}^\delta V_\delta$ where V_δ is that portion of the compactification volume which produces power-law running for α_{GUT} .

Thus, to summarize, we have shown that it is possible for “GUT precursor” states to appear with masses that are extremely light compared with the scale of gauge coupling unification. This suggests a possible new TeV-scale direction for orbifold GUT models. Indeed, more generally, we have seen that ultraviolet embeddings into fixed-point theories can be used to provide a new method for maintaining or stabilizing a wide separation of energy scales within a single model. Using this technique, “hybrid” models with coexisting high and low energy scales can therefore be constructed in a variety of contexts. Equally importantly, however, our four-dimensional “Kaluza-Klein” realization of such fixed points should also provide a new technique for the study of such theories and their properties under various compactifications, both with and without supersymmetry breaking and gauge symmetry breaking. These and other directions await exploration.

ACKNOWLEDGMENTS

KRD was supported in part by the National Science Foundation under Grant PHY-0071054, and by a Research Innovation Award from Research Corporation. ED was supported in part by the RTN European Program HPRN-CT-2000-00148. We wish to thank C. Grojean, K. Intriligator, G.D. Kribs, J. Mourad, M.J. Strassler and especially E. Poppitz for discussions. KRD and TG wish to acknowledge the hospitality of the Aspen Center for Physics where this work was initiated, and KRD and ED wish to acknowledge the hospitality of the Theoretical Physics Institute at the University of Minnesota where this work was completed.

-
- [1] I. Antoniadis, Phys. Lett. B **246** (1990) 377; I. Antoniadis, K. Benakli and M. Quiros, Phys. Lett. B **331** (1994) 313 [arXiv:hep-ph/9403290].
 - [2] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **429** (1998) 263 [arXiv:hep-ph/9803315]; I. Antoniadis *et al.*, Phys. Lett. B **436** (1998) 257 [arXiv:hep-ph/9804398].
 - [3] K. R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. B **436** (1998) 55 [arXiv:hep-ph/9803466]; Nucl. Phys. B **537** (1999) 47 [arXiv:hep-ph/9806292]; arXiv:hep-ph/9807522.
 - [4] E. Witten, Nucl. Phys. B **471** (1996) 135 [arXiv:hep-th/9602070]; J. D. Lykken, Phys. Rev. D **54** (1996) 3693 [arXiv:hep-th/9603133]; G. Shiu and S.-H. H. Tye,

- Phys. Rev. D **58** (1998) 106007 [arXiv:hep-th/9805157]; Z. Kakushadze and S.-H. H. Tye, Nucl. Phys. B **548** (1999) 180 [arXiv:hep-th/9809147].
- [5] L. J. Dixon, J. A. Harvey, C. Vafa and E. Witten, Nucl. Phys. B **261** (1985) 678; Nucl. Phys. B **274** (1986) 285.
- [6] See, *e.g.*, M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory* (Cambridge University Press, 1987); J. Polchinski, *String Theory* (Cambridge University Press, 1998).
- [7] See, *e.g.*, I. Antoniadis, J. R. Ellis, J. S. Hagelin and D. V. Nanopoulos, Phys. Lett. B **231** (1989) 65; A. E. Faraggi, Phys. Lett. B **274** (1992) 47; Phys. Lett. B **278** (1992) 131; Phys. Lett. B **520** (2001) 337 [arXiv:hep-ph/0107094].
- [8] For phenomenological reviews, see, *e.g.*, L. E. Ibáñez, arXiv:hep-th/9112050; arXiv:hep-th/9505098; I. Antoniadis, arXiv:hep-th/9307002; M. Dine, arXiv:hep-ph/9309319; A. E. Faraggi, Nucl. Phys. B **387** (1992) 239 [arXiv:hep-th/9208024]; arXiv:hep-ph/9405357; K. R. Dienes and A. E. Faraggi, Phys. Rev. Lett. **75** (1995) 2646 [arXiv:hep-th/9505018]; Nucl. Phys. B **457** (1995) 409 [arXiv:hep-th/9505046]; K. R. Dienes, A. E. Faraggi and J. March-Russell, Nucl. Phys. B **467** (1996) 44 [arXiv:hep-th/9510223]; J. L. Lopez and D. V. Nanopoulos, arXiv:hep-ph/9511266; J. D. Lykken, arXiv:hep-ph/9511456; Z. Kakushadze and S.-H. H. Tye, arXiv:hep-th/9512155; F. Quevedo, arXiv:hep-th/9603074; arXiv:hep-ph/9707434; K. R. Dienes, Phys. Rept. **287** (1997) 447 [arXiv:hep-th/9602045]; Z. Kakushadze, G. Shiu, S.-H.H. Tye and Y. Vtorov-Karevsky, Int. J. Mod. Phys. **13** (1998) 2551.
- [9] Y. Kawamura, Prog. Theor. Phys. **105** (2001) 999 [arXiv:hep-ph/0012125]; Prog. Theor. Phys. **105** (2001) 691 [arXiv:hep-ph/0012352]; T. Kawamoto and Y. Kawamura, arXiv:hep-ph/0106163.
- [10] See, *e.g.*, L. J. Hall and Y. Nomura, Phys. Rev. D **64** (2001) 055003 [arXiv:hep-ph/0103125]; Phys. Rev. D **65** (2002) 125012 [arXiv:hep-ph/0111068]; arXiv:hep-ph/0205067; arXiv:hep-ph/0207079; Y. Nomura, D. R. Smith and N. Weiner, Nucl. Phys. B **613** (2001) 147 [arXiv:hep-ph/0104041]; R. Barbieri, L. J. Hall and Y. Nomura, Phys. Rev. D **66** (2002) 045025 [arXiv:hep-ph/0106190]; L. J. Hall, Y. Nomura and D. R. Smith, Nucl. Phys. B **639** (2002) 307 [arXiv:hep-ph/0107331]; L. J. Hall, Y. Nomura, T. Okui and D. R. Smith, Phys. Rev. D **65** (2002) 035008 [arXiv:hep-ph/0108071].
- [11] See, *e.g.*, A. Hebecker and J. March-Russell, Nucl. Phys. B **613** (2001) 3 [arXiv:hep-ph/0106166]; Nucl. Phys. B **625** (2002) 128 [arXiv:hep-ph/0107039]; Phys. Lett. B **539** (2002) 119 [arXiv:hep-ph/0204037]; Phys. Lett. B **541** (2002) 338 [arXiv:hep-ph/0205143]; L. Hall, J. March-Russell, T. Okui and D. R. Smith, arXiv:hep-ph/0108161.
- [12] See, *e.g.*, G. Altarelli and F. Feruglio, Phys. Lett. B **511** (2001) 257 [arXiv:hep-ph/0102301]; J. A. Bagger, F. Feruglio and F. Zwirner, Phys. Rev. Lett. **88** (2002) 101601 [arXiv:hep-th/0107128]; T. j. Li, Phys. Lett. B **520** (2001) 377 [arXiv:hep-th/0107136]; Nucl. Phys. B **619** (2001) 75 [arXiv:hep-ph/0108120]; Nucl. Phys. B **633** (2002) 83 [arXiv:hep-th/0112255]; N. Haba *et al.*, Prog. Theor. Phys. **106** (2001) 1247 [arXiv:hep-ph/0108003]; T. Asaka, W. Buchmuller and L. Covi, Phys. Lett. B **523** (2001) 199 [arXiv:hep-ph/0108021]; R. Dermisek and A. Mafi, Phys. Rev. D **65** (2002) 055002 [arXiv:hep-ph/0108139]; C. S. Huang *et al.*, Phys. Lett. B **530** (2002) 218 [arXiv:hep-th/0112046]; N. Haba, T. Kondo and Y. Shimizu, Phys. Lett. B **531** (2002) 245 [arXiv:hep-ph/0112132].
- [13] T. R. Taylor and G. Veneziano, Phys. Lett. B **212** (1988) 147.
- [14] K. Agashe, JHEP **0105** (2001) 017 [arXiv:hep-ph/0012182].
- [15] Z. Kakushadze and T. R. Taylor, Nucl. Phys. B **562** (1999) 78 [arXiv:hep-th/9905137].
- [16] R. Contino, L. Pilo, R. Rattazzi and E. Trincherini, Nucl. Phys. B **622** (2002) 227 [arXiv:hep-ph/0108102].
- [17] N. Seiberg, Phys. Lett. B **388** (1996) 753 [arXiv:hep-th/9608111]; K. A. Intriligator, D. R. Morrison and N. Seiberg, Nucl. Phys. B **497** (1997) 56 [arXiv:hep-th/9702198]; D. I. Kazakov, arXiv:hep-th/0209100.
- [18] N. Seiberg, Phys. Lett. B **390** (1997) 169 [arXiv:hep-th/9609161].
- [19] N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Rev. Lett. **86** (2001) 4757 [arXiv:hep-th/0104005]; C. T. Hill, S. Pokorski and J. Wang, Phys. Rev. D **64** (2001) 105005 [arXiv:hep-th/0104035].
- [20] C. Csaki, J. Erlich, C. Grojean and G. D. Kribs, Phys. Rev. D **65** (2002) 015003 [arXiv:hep-ph/0106044].
- [21] N. Seiberg, Nucl. Phys. B **435** (1995) 129 [arXiv:hep-th/9411149].
- [22] V. A. Novikov, M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **229** (1983) 381; M. A. Shifman and A. I. Vainshtein, Nucl. Phys. B **277** (1986) 456; Nucl. Phys. B **359** (1991) 571.
- [23] We thank K. Agashe, A. Delgado and Y. Nomura for discussions of this issue [private communication].
- [24] K. R. Dienes, E. Dudas and T. Gherghetta, November 2002, as reported in a talk by E. D. at PAS-COS '03, January 2003. Scanned transparencies available at <http://theory.theory.tifr.res.in/pascos>.
- [25] See, *e.g.*, F. P. Correia, M. G. Schmidt and Z. Tavartkiladze, hep-ph/0302038.
- [26] N. Arkani-Hamed and M. Schmaltz, Phys. Rev. D **61** (2000) 033005 [arXiv:hep-ph/9903417].
- [27] Z. Kakushadze, Nucl. Phys. B **552** (1999) 3 [arXiv:hep-th/9812163].
- [28] See, *e.g.*, R. Sundrum, Phys. Rev. D **59** (1999) 085010 [arXiv:hep-ph/9807348]; N. Arkani-Hamed, S. Dimopoulos and J. March-Russell, Phys. Rev. D **63** (2001) 064020 [arXiv:hep-th/9809124]; K. R. Dienes, E. Dudas, T. Gherghetta and A. Riotto, Nucl. Phys. B **543** (1999) 387 [arXiv:hep-ph/9809406]; E. Ponton and E. Poppitz, JHEP **0106** (2001) 019 [arXiv:hep-ph/0105021].
- [29] K.R. Dienes, E. Dudas, T. Gherghetta and E. Poppitz, in progress.
- [30] P. H. Frampton and C. Vafa, arXiv:hep-th/9903226.
- [31] See, *e.g.*, A. Pomarol, Phys. Rev. Lett. **85** (2000) 4004 [arXiv:hep-ph/0005293]; W. D. Goldberger, Y. Nomura and D. R. Smith, arXiv:hep-ph/0209158.
- [32] K. R. Dienes, Nucl. Phys. B **611** (2001) 146 [arXiv:hep-ph/0104274].